

POST SURGERY INSTRUCTIONS FOR FUE

Dear Patient,

Following your surgery procedure, special care must be taken especially during the first two weeks.

Please read the below instructions carefully:

- The use of the bandana should be limited for getting around in public. Do not leave on for many hours at a time and especially do not sleep with it on.
- Direct trauma is one of the main causes of graft loss after transplantation so take care to avoid bumping your head especially when getting in and out of the car.

First 3 Nights

Sleep with 3-4 pillows behind your shoulders in order to keep your head above your heart, approximately at a 45 degree angle. This is important as it will minimize swelling.

Report immediately to clinic

- Temperature above 38° C
- Any excess bleeding
- Sudden swelling within first 12 hours of surgery

1. Care of Recipient and Donor Areas

1st Day The day following your transplant you must spray the transplanted (recipient) area with saline solution every hour and must continue doing so until it runs out. Apply Aloe Vera Gel mixed with Fucicort and Vitamin E capsules to donor area twice daily. See below for details on mixing.

2nd-5th Day Wet scalp with lukewarm water WITHOUT direct shower pressure to the scalp. Apply **Stellisept Scrub** to hands and work up a rich lather. Apply to all areas of the scalp with a gentle touch but DO NOT rub in, dab gently. Make very light circular motions with fingertips and then leave to soak for 5 minutes. Rinse, once again taking care that the water does NOT run with direct shower pressure onto the grafts. You can use your hand to block the shower head to achieve reduced water pressure. Repeat but this time using **After Transplant Shampoo**. Apply **Aloe Vera Gel mixed with Fucicort and Vitamin E capsules**. This whole procedure should be done twice daily.

6th-13th Day Wet the scalp with lukewarm water without excessive direct shower pressure to the grafts. Apply **After Transplant shampoo** to hands and work up a rich lather. Apply to all areas of the scalp and use your finger tips to gently massage (with circular motions) for 3 minutes. Leave to soak for another 5 minutes and then massage again for another 3 minutes before rinsing. This will help to soften up the scabs which will subsequently clear up. Rinse without direct shower pressure. This whole procedure should be done twice daily until scabs have cleared up.

Day 14th and onwards You may now use direct shower pressure. Continue using After Transplant Shampoo until it is finished and then you can use your regular shampoo or a shampoo recommended by the clinic.

HDC Medical Trichology Centre

Protogoras 6, Ayios Antonios
1505 Nicosia
Cyprus

Tel: (+357) 22 345 245
Fax: (+357) 22 349 348
Email: info@hdc.com.cy

► Important ◀

Avoid any kind of cream, foam, gel or any other hair styling products for at least 15 days after the transplantation. Hair dyes should not be used for at least 45 days post operatively and with the consent of the clinic.

**Instructions
on mixing
Aloe Vera
with
Fucicort
and Vitamin
E capsules**

You will need to mix this up every time you apply it, as it must be fresh. This mixture will help the healing process and will minimize any scarring. It is important that you use twice daily from the day following your surgery (day 1) until it runs out. You will mix up in the following way:

1. Fill up the base of your small cup with Aloe Vera Gel so that it covers the base when it is spread out.
2. Add in about 2cm of Fucicort cream
3. Add in four Vitamin E capsules by piercing them with a needle and squeezing into the mixture.
4. Mix with spatula and apply to donor area and leave to absorb.

Scalp drops

Scalp drops plus 2 are to be used from the 10th day after surgery. It is recommended that they are continued for at least 3 months to help minimize shock loss. Basically scalp drops should be applied to a clean dry scalp twice daily.

1. Shake bottle
2. Draw drops half way up the pipette (This will make 1ml)
3. Apply to top part of scalp slowly with pipette and use pipette to spread.
4. Leave to absorb on its own. Do not massage.
5. Do not worry if you have not covered whole top area with your 1ml. There will be local absorption so all areas will be covered.

► Important ◀

If you have an allergic reaction to the drops stop using them and contact our Trichologist Stella Prokopiou for further advice.

3. Medication

**Sleeping
Pill**

You will be given a pill that will help you sleep the first night after the transplantation. Do not drive or handle heavy machinery after taking this pill. The consumption of alcohol is also forbidden whilst you are taking medication.

Pain Killers

Our aim is to reduce any pain and discomfort following your transplantation. You will be given pain killers which you can take every six hours. It is forbidden for you to take aspirin or any anti inflammatory medication for at least three days after the transplant in order to avoid any bleeding.

4. Additional Useful Information

HDC Medical Trichology Centre

Protogoras 6, Ayios Antonios
1505 Nicosia
Cyprus

Tel: (+357) 22 345 245
Fax: (+357) 22 349 348
Email: info@hdc.com.cy

Hiccups It is advisable to avoid drinking any carbonated drink (soft drink, beer) for 72 hours after the transplant. By this, you will have a reduced possibility of developing hiccups which can result in damage to the transplanted area. In the event that you develop hiccups, please call the HDC Centre.

Blood loss The loss of blood will be noticed immediately following your transplant procedure and while at the Centre. It is rare to experience blood loss once you have left the centre. In such an instance, apply continuous pressure on the point where bleeding is occurring with a clean cloth for 20 minutes. In the event that bleeding persists, please contact the Centre immediately.

Rest Position The ideal position to place your head for the first 3 nights is a 45° angle to the bed. You can use as many pillows as you like for support. This position reduces the possibilities of any swelling.

Swelling You will be given medication for the reduction of swelling on the transplanted area. If swelling appears after the end of the transplant process, this reduces on its own within 3-4 days and is harmless. If it is painful and/or is accompanied by fever then contact the Centre immediately.

Exercise Alcohol Smoking The following day from your transplant procedure, it is advisable that you stay home and rest. The following three days you can begin taking on light, day to day work. At the same time, you should avoid any form of exercise for the first 15 days, especially lifting of heavy objects. After 15 days light exercise eg. Use of treadmill or swimming is okay. Heavy exercise like weight lifting should be avoided during the first month. Avoid any form of sexual contact for at least the first three days. We also advise no smoking or alcohol for the first three days following your procedure as this can lessen the growth of transplants.

Sun Please avoid unprotected sunlight for at least three months, this is because you might get sunburn without feeling it. Please use a baseball cap and apply a strong sunscreen cream over SPF 35

Infection On very rare occasions, a transplant may become infected. Slight skin irritation, swelling and discomfort may appear in the first days. If these symptoms persist and even increase after the third day, please contact the Centre immediately.

Numbness In the very unlikely event that you develop **numbness**, do not worry. This is due to the severing of small nerves during the transplant procedure. Usually the numbness, if present, will fade away over a period of two months, as the nerves repair themselves.



For any questions or queries, please do not hesitate to
Contact us at 777 777 19